

Annual Nutrient Loadings, Primary Productivity, and Trophic State of Lake Koocanusa, Montana and British Columbia, 1972–80

GEOLOGICAL SURVEY PROFESSIONAL PAPER 1283

*Prepared for the U.S. Army Corps of
Engineers, Seattle District*



Annual Nutrient Loadings, Primary Productivity, and Trophic State of Lake Koocanusa, Montana and British Columbia, 1972–80

By PAUL F. WOODS

GEOLOGICAL SURVEY PROFESSIONAL PAPER 1283

*Prepared for the U.S. Army Corps of
Engineers, Seattle District*



UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON : 1982

UNITED STATES DEPARTMENT OF THE INTERIOR

JAMES G. WATT, *Secretary*

GEOLOGICAL SURVEY

Dallas L. Peck, *Director*

Library of Congress Cataloging in Publication Data

Woods, Paul F.

Annual nutrient loadings, primary productivity, and trophic state of Lake Koochanusa, Montana and British Columbia, 1972-80.

(Geological Survey Professional Paper 1283)

Bibliography: 24 p.

Supt. of Docs. No.: I 19.16: 1283

1. Primary productivity (Biology)—Koochanusa, Lake (B.C. and Mont.) 2. Eutrophication—Koochanusa, Lake (B.C. and Mont.) 3. Water quality—Koochanusa, Lake (B.C. and Mont.) 4. Koochanusa, Lake (B.C. and Mont.)

I. United States Army Corps of Engineers. Seattle, Washington, District. II. Title. III. Series.

QH105.M9W66 1982 574 82-600264

CONTENTS

	Page
Abstract	1
Introduction	1
Regional setting	3
Hydrology	4
Nutrient loadings	5
Methodology	6
Influent loadings	7
Loadings discharged from Lake Kooconusa	8
Nutrient retention in Lake Kooconusa	8
Primary productivity	10
Methodology	10
Seasonal trend	10
Annual trend	12
Relationship of annual areal primary productivity and selected environmental variables	12
Chlorophyll <i>a</i> distribution within the water column	18
Trophic state of Lake Kooconusa	20
Conclusions	22
References	24

ILLUSTRATIONS

	Page
FIGURE 1. Map showing location of Kootenai River drainage basin and Lake Kooconusa	2
2-4. Graphs showing:	
2. Monthly and annual inflow to and releases from Lake Kooconusa, 1972-80	5
3. Daily variation in surface altitude of Lake Kooconusa, 1972-80	6
4. Daily values of areal primary productivity at the four limnological stations, Lake Kooconusa, 1972-80	13
5. Scatterplots of annual areal primary productivity versus maximum and mean annual euphotic zone depths, annual areal phosphorus loading, and annual nitrogen retention coefficient for Lake Kooconusa, 1972-80	16
6. Graph showing annual variation of mean and maximum euphotic zone depths, areal primary productivity, areal phosphorus loading, and nitrogen retention coefficient for Lake Kooconusa, 1972-80	17
7. Typical diagram used to quantify the percentage distribution of chlorophyll <i>a</i> within the water column of Lake Kooconusa.	19
8. Graph showing application of Lake Kooconusa data in table 17 to Vollenweider total phosphorus loading and mean depth/hydraulic-residence time diagram	22
9. Graph showing application of Lake Kooconusa data in table 17 to Vollenweider total nitrogen loading and mean depth/hydraulic-residence time diagram	23

TABLES

	Page
TABLE 1. Annual precipitation at climatological station 1 and annual mean air temperature at climatological station 2, 1972–80	3
2. Volume and surface area of Lake Koochanusa at selected reservoir surface altitudes	4
3. Lake-filling and hydraulic-residence times of Lake Koochanusa, 1972–80	6
4. Annual loadings of total phosphorus that entered Lake Koochanusa, 1970–80	7
5. Annual loadings of total nitrogen that entered Lake Koochanusa, 1970–80	8
6. Comparison of 1973–78 loadings of total phosphorus (TP) and total nitrogen (TN) at two stations on the St. Mary River, one upstream and one downstream from the fertilizer plant at Kimberley, British Columbia	8
7. Annual loadings of total phosphorus (TP) and total nitrogen (TN) discharged from Lake Koochanusa, 1970–80	9
8. Nutrient-retention coefficients for total phosphorus and total nitrogen in Lake Koochanusa, 1972–80	9
9. Statistical summary of daily areal primary productivity and euphotic zone depth at the four limnological stations, Lake Koochanusa, 1972–80	11
10. Statistical summary of daily areal primary productivity in Lake Koochanusa, 1972–80	12
11. Annual areal primary productivity in Lake Koochanusa, 1972–80	12
12. Statistical summary of environmental variables used in correlation and multiple regression analyses of annual areal primary productivity in Lake Koochanusa, 1972–80	14
13. Correlation coefficients between annual areal primary productivity and environmental variables X1 through X12	14
14. Multiple regression models for prediction of annual areal primary productivity in Lake Koochanusa	15
15. Multiple regression model developed by Woods (1979) for prediction of daily areal primary productivity in Lake Koochanusa	18
16. Percentage distribution of chlorophyll <i>a</i> in three depth strata at the Forebay and Tenmile stations in 1980	20
17. Required data for using Vollenweider nutrient loading models to determine the trophic state of Lake Koochanusa	21

METRIC CONVERSION TABLE

To convert International System (SI) of units in this report to inch-pound units, multiply by the following factors:

<i>Multiply SI unit</i>	<i>By</i>	<i>To obtain inch-pound unit</i>
cubic kilometer (km ³)	8.107 × 10 ⁵	acre-foot
cubic meter per second (m ³ ·s ⁻¹)	3.531 × 10 ¹	cubic foot per second
gram (g)	2.205 × 10 ⁻³	pound
gram per square centimeter (g·cm ⁻²)	2.049	pound per square foot
gram per square meter (g·m ⁻²)	2.049 × 10 ⁻⁴	pound per square foot
gram per square meter per year (g·m ⁻² ·a ⁻¹)	2.049 × 10 ⁻⁴	pound per square foot per year
hectare (ha)	2.471	acre
kilogram per hectare per year (kg·ha ⁻¹ ·a ⁻¹)	8.922 × 10 ⁻¹	pound per acre per year
kilometer (km)	6.214 × 10 ⁻¹	mile
lux (lx)	9.290 × 10 ⁻²	foot-candle
meter (m)	3.281	foot
megagram (Mg)	1.102	ton (short)
megagram per cubic kilometer (Mg·km ⁻³)	1.359 × 10 ⁻⁶	ton (short) per acre-foot
milligram (mg)	2.205 × 10 ⁻⁶	pound
milligram per cubic meter (mg·m ⁻³)	6.245 × 10 ⁻⁸	pound per cubic foot
milligram per square meter per day (mg·m ⁻² ·d ⁻¹)	2.049	pound per square foot per day
millimeter (mm)	3.937 × 10 ⁻²	inch
square kilometer (km ²)	3.861 × 10 ⁻¹	square mile

Temperature in degrees Celsius (°C) can be converted to degrees Fahrenheit (°F) by the following equation:

$$^{\circ}\text{F} = 1.8 \times ^{\circ}\text{C} + 32$$

National Geodetic Vertical Datum of 1929 (NGVD of 1929): A geodetic datum derived from a general adjustment of the first-order level nets of both the United States and Canada, formerly called "mean sea level." NGVD of 1929 is referred to as sea level in this report.

ANNUAL NUTRIENT LOADINGS, PRIMARY PRODUCTIVITY, AND TROPHIC STATE OF LAKE KOOCANUSA, MONTANA AND BRITISH COLUMBIA, 1972-80

By PAUL F. WOODS

ABSTRACT

Limnological data collected at Lake Koocanusa were used to investigate the relationship of nutrient loadings, primary productivity, and trophic state of the reservoir during 1972-80. The reservoir, on the Kootenai River, was impounded by Libby Dam on March 21, 1972. Manipulation of the 7.16-cubic-kilometer reservoir for flood control, its primary function, created large fluctuations in reservoir volume and produced annual lake-filling times that ranged from 0.14 to 0.66 year.

Loadings of nitrogen and phosphorus prior to and following impoundment of Lake Koocanusa were found to be large enough to predict eutrophic conditions. Beginning in 1976, total phosphorus loadings, but not total nitrogen loadings, were substantially reduced following improvements in waste-water treatment at a fertilizer plant located upstream from the reservoir. The closure of Libby Dam substantially reduced loadings of nitrogen and phosphorus downstream from Lake Koocanusa. On the average, the reservoir retained 63 percent of its influent loading of total phosphorus and 25 percent of its influent loading of total nitrogen.

Daily areal and volumetric primary productivity varied widely in each year at four sampled limnological stations. During the 9 years studied, daily areal primary productivity, in milligrams of carbon fixed per square meter, ranged from 0.4 to 420.0; the mean of the 313 sampled days was 128.5. Annual areal primary productivity ranged from 23.2 to 38.5 grams of carbon fixed per square meter and thereby categorized Lake Koocanusa as oligotrophic.

The relationship of annual areal primary productivity and 12 selected environmental variables was determined by multiple regression analysis. One of the models that was derived used two variables—annual euphotic zone depth and annual areal phosphorus loading—and accounted for 62.0 percent of the variation in annual areal primary productivity.

The distribution of chlorophyll *a* within the water column indicated that, on the average, more than one-half of the phytoplankton in the reservoir was beneath the euphotic zone. These results support the hypothesis that the reservoir's weak thermal structure had allowed circulation of phytoplankton out of the euphotic zone.

The trophic state of Lake Koocanusa was categorized as eutrophic when based on the relationship of the nutrient loadings and the reservoir's ratio of mean depth to hydraulic-residence time. This result conflicted with the oligotrophic ranking the reservoir received based on its areal primary productivity. The discrepancy in trophic state was attributed mainly to the failure of nutrient loading models to adequately account for physical processes within reservoirs. Part of the nutrient loading that entered Lake Koocanusa was unavailable to phytoplankton because the nutrients were carried beneath the euphotic zone by large volumes of interflow and underflow. Another

part of the nutrient loading was adsorbed to suspended sediment and removed from the water column. Thus, phytoplankton primary productivity was controlled not only by nutrients, but also by other limnological processes.

INTRODUCTION

Libby Dam was constructed as part of a treaty between the United States and Canada to cooperatively develop the water resources of the Columbia River drainage basin. Construction of the dam on the Kootenai (spelled Kootenay in Canada) River began in 1966, and Lake Koocanusa was officially impounded on March 21, 1972. The reservoir has a volume of 7.16 km³ and provides flood storage, hydroelectric power production, and recreation benefits. The 148-km-long reservoir straddles the United States-Canadian border and impounds water from about 23,271 km², or 47 percent, of the Kootenai River drainage basin (fig. 1). Three Canadian rivers, the Kootenai, Elk, and Bull, supply 87 percent of the reservoir's inflow (Bonde, 1979) and, therefore, exert a major influence on the limnology of Lake Koocanusa.

Preimpoundment water-quality studies indicated the presence of large concentrations of total phosphorus, orthophosphorus¹, and total nitrogen in the Kootenai River, which were attributed to industrial point-source discharges in the Canadian part of the drainage basin (Bonde and Bush, 1975). Using the preimpoundment data, these authors calculated that the reservoir's surface would receive areal loadings of 10 g·m⁻² of total phosphorus and 20 g·m⁻² of total nitrogen. Areal nutrient loadings have been used in conjunction with mean depth and hydraulic residence time to estimate a water body's susceptibility to eutrophication (Vollenweider,

¹Rigler (1973) reported that methodological problems preclude determination of the actual concentration of orthophosphate-phosphorus in natural water samples. He suggested use of the term "soluble reactive phosphorus" as a more accurate operational definition. However, the term orthophosphorus is retained for use in this report because it is the terminology currently used by the Geological Survey.

1968, 1975). Application of Vollenweider's methods to Lake Koocanusa by Bonde and Bush (1975) indicated that annual areal loadings of $2.0 \text{ g}\cdot\text{m}^{-2}$ of total phospho-

rus and $8.0 \text{ g}\cdot\text{m}^{-2}$ of total nitrogen would be sufficient to cause concern for eutrophication of the reservoir. These values were substantially less than the predicted




FIGURE 1.—Location of Kootenai River drainage basin and Lake Koocanusa.

annual loadings; therefore, Bonde and Bush considered Lake Kooconusa to be susceptible to eutrophication. However, they cautioned that prediction of the actual response of the reservoir to such nutrient loadings would be difficult.

Bonde and Bush (1975) reported that blooms of *Aphanizomenon flos-aquae*, a blue-green alga characteristic of eutrophic waters, occurred in Lake Kooconusa in the autumns of 1974 and 1975 and considered these symptomatic evidence of eutrophication. However, Woods (1979) reported that annual primary productivity in the reservoir during 1972 through 1975 was characteristic of oligotrophic, not eutrophic, water bodies. Preliminary evaluations by Hobbie (1976) and Bush and Bonde (1977) revealed that primary productivity in Lake Kooconusa was predominantly controlled by physical limnological processes. Additional studies by Woods (1979, 1981) attributed the reservoir's oligotrophy to suppression of phytoplankton photosynthesis by circulation of phytoplankton out of the euphotic zone.

Subsequent to the 1972-75 period evaluated by Woods (1979), waste-water treatment was improved at the industrial point-source discharges in Canada. Beginning in 1976, these improvements resulted in substantial reductions in loadings of total phosphorus delivered into Lake Kooconusa (Bonde, 1979).

Although the limnological data collected during 1972 through 1975 have been evaluated (Woods, 1979, 1981; Woods and Falter, 1982), little such work has been done on the post-1975 data. The U.S. Army Corps of Engineers, therefore, requested that the U.S. Geological Survey further evaluate some selected data. The Corps of Engineers was primarily interested in furthering their understanding of the limnological processes that control the trophic state of Lake Kooconusa. Annual nutrient loadings and annual areal primary productivity were specified for intensive evaluation, because both have been extensively used to classify the trophic state of lakes and reservoirs.

The purpose of this resultant report is to categorize the trophic state of Lake Kooconusa based on annual nutrient loadings and annual areal primary productivity. Seasonal and annual trends in primary productivity are described, as well as annual trends in loadings of nitrogen and phosphorus. Control of annual areal primary productivity by selected limnological processes in Lake Kooconusa is evaluated in relation to results derived from multiple regression analyses and an analysis of the distribution of chlorophyll *a* within the water column. The report addresses the 1972-80 postimpoundment phase and is based on preimpoundment data collected after 1967 and postimpoundment data collected from 1972 through 1980.

REGIONAL SETTING

Lake Kooconusa lies within the Kootenai River drainage basin, an area of 49,987 km² occupying parts of Idaho, Montana, and British Columbia. Altitudes in the basin range from 3,618 m above sea level at Mt. Assiniboine to 418 m at the confluence of the Kootenai and Columbia Rivers near Castlegar, British Columbia. The surface of Lake Kooconusa at maximum pool is at an altitude of 749.50 m.

The Kootenai River, the principal tributary to Lake Kooconusa, arises 64 km west of Banff, Alberta, and flows southward to the Purcell Mountains. Near the Purcells, the river is impounded by Libby Dam 4.5 km upstream from the Fisher River. The Kootenai River eventually enters Kootenay Lake, and thence the Columbia River near Castlegar, British Columbia.

Between Canal Flats, British Columbia, and the Tobacco River, the Kootenai River flows along the Rocky Mountain Trench—a graben having a floor about 11 km wide and a vertical displacement of several hundred meters. The trench is partly filled with Pleistocene and Holocene deposits of unconsolidated silt, sand, and gravel into which the Kootenai River has incised about 60 to 90 m (Coffin, 1970). Except for the relatively flat terrain of the Rocky Mountain Trench, the topography upstream from Libby Dam is dominated by rugged mountains that are densely forested with conifers.

Climatically, the Kootenai River basin is subjected to maritime influences in the winter and continental influences in the summer. These influences, in conjunction with the mountainous terrain, result in complex weather patterns. The Rocky Mountain Trench receives

TABLE 1.—Annual precipitation at climatological station 1^a and annual mean air temperature at climatological station 2^b, 1972-80

Year	Annual precipitation, in millimeters		Annual mean air temperature, in degrees Celsius
	Total	Departure from normal ¹	
1972	485.6	-7.2	6.2
1973	357.4	-135.4	6.6
1974	460.2	-32.6	7.4
1975	527.6	+34.8	6.1
1976	354.3	-138.5	7.0
1977	403.1	-89.7	7.2
1978	424.4	-68.4	6.2
1979	311.7	-181.1	7.1
1980	551.9	+59.1	7.1

^aClimatological station 1 is "Libby 1 NE Ranger Station" in reports of the U.S. Department of Commerce (issued annually).

^bClimatological station 2 is "Libby Dam" in reports of the U.S. Department of Commerce (issued annually).

^cDeparture from long-term annual average precipitation of 492.8 mm for 83 years of record.

an average of 400 mm of precipitation per year and is considered semiarid, whereas the Rocky Mountains annually receive an average of 1,000 mm of precipitation (Water Resources Service, 1976). About 70 percent of the basin's precipitation falls as snow during November through March (Bonde and Bush, 1975). At climatological station 1 (fig. 1), precipitation has been recorded for 83 years and temperature for 73 years (U.S. Department of Commerce, issued annually). Based on these data, the mean annual precipitation near Libby Dam is 492.8 mm and the mean annual temperature is 7.3°C.

Annual precipitation and annual mean air temperatures recorded at climatological stations 1 and 2 (fig. 1) near Libby Dam during 1972-80 are listed in table 1. Precipitation ranged from 311.7 to 551.9 mm. Except for 1975 and 1980, precipitation was less than normal over the 9 years, particularly in 1973, 1976, and 1979. Yearly variation in air temperature was small, ranging from 6.1° to 7.4°C, but the monthly values published by the U.S. Department of Commerce (issued annually) varied widely.

HYDROLOGY

The predominant flood-control function of Lake Koocanusa necessitates substantial reductions in volume during the autumn and winter to provide adequate capacity for spring snowmelt runoff. This operational schedule produces a wide range of reservoir volume and surface area (table 2). At maximum pool, the reservoir contains 7.16 km³ and has a surface area of about 188 km². A drawdown of 52.4 m to minimum operational pool reduces the volume to 1.08 km³ and the surface

TABLE 2.—Volume and surface area of Lake Koocanusa at selected reservoir surface altitudes

Surface altitude, in meters	Volume, in cubic kilometers	Percent of maximum volume	Surface area, in square kilometers	Percent of maximum area
^a 749.50	7.16	100.0	188.2	100.0
740.00	5.52	77.1	159.4	84.7
730.00	4.08	57.0	131.5	69.9
720.00	2.91	40.6	103.2	54.8
710.00	1.99	27.8	80.9	43.0
700.00	1.26	17.6	64.8	34.4
^b 697.08	1.08	15.1	59.1	31.4
690.00	.71	9.9	46.5	24.7
680.00	.34	4.7	27.5	14.6
^c 671.17	.14	2.0	16.2	8.6

^aMaximum pool.

^bMinimum operational pool.

^cDead storage.

area to 59.1 km².

Annual inflow to the reservoir exceeded or equaled annual outflow from Libby Dam except in 1977 and 1979, the 2 years in which inflow was about one-half of normal (fig. 2). Maximum inflow occurred in May or June, whereas, in general, outflow was largest during reservoir drawdown. This cyclic pattern of inflow and outflow produced large fluctuations in reservoir volume and surface altitude (fig. 3). The reservoir surface generally was at its yearly maximum altitude in July, August, and September and at its minimum during March and April.

The reservoir attained its maximum surface altitude for less than 7 months during the 9 years graphed in figure 3. Three months was the longest period in which the reservoir surface was at a constant altitude. Most of the 9 years was characterized by a constantly changing reservoir surface altitude and, hence, volume.

Reservoir volume, inflow, and outflow can be related mathematically as lake-filling time and hydraulic-residence time. Lake-filling time represents the time required to replace the volume of a reservoir at a given inflow (Daley and others, 1981); whereas hydraulic-residence time represents the time required to replace the volume of a reservoir at a given outflow (Vollenweider, 1975). The two are calculated by the following equations:

$$R = \frac{V}{I} \quad (1)$$

$$F = \frac{V}{O} \quad (2)$$

where

R is lake-filling time, in years;

F is hydraulic-residence time, in years;

V is reservoir volume, in cubic kilometers;

I is annual inflow, in cubic kilometers; and

O is annual outflow, in cubic kilometers.

The theoretical lake-filling time of Lake Koocanusa is 0.67 year, based on a reservoir volume of 7.16 km³ and an assumed annual inflow of 10.65 km³. However, the reservoir experienced a large range in volume, inflow, and outflow from 1972 through 1980, which yielded annual lake-filling times that ranged from 0.14 to 0.66 year and annual hydraulic-residence times that ranged from 0.14 to 0.62 year (table 3). In addition to the annual variation, the reservoir exhibited monthly variation in volume, inflow, and outflow. Lake-filling and hydraulic-residence times, therefore, were calculated using monthly data instead of annual data. The results, in table 3, indicated a wide range in values. For example, the mean monthly lake filling time in 1976 was 0.71 year, but during that year, the value ranged from 0.13 to 1.54 years. The minimum value resulted




FIGURE 2.—Monthly and annual inflow to and releases from Lake Koochanusa, 1972–80.

from springtime snowmelt runoff entering a partly drawdown reservoir in May; the maximum lake-filling time of 1.54 years occurred in December as small inflow entered a large reservoir volume.

NUTRIENT LOADINGS

The relationship between nutrient loadings and the trophic state of a lake or reservoir has been extensively studied and was reviewed in recent reports by Rast and Lee (1978) and Reckhow (1979). A model developed by Vollenweider (1968, 1975) was discussed in these two reports. This model, based on areal loadings of total phosphorus or total nitrogen, mean depth, and hydraulic-residence time, plots a point onto a graph from which the trophic state of the lake or reservoir may be estimated. In Lake Koochanusa, loadings of both nutrients

were large enough to produce a eutrophic ranking (Bonde and Bush, 1975).

The source of much of the phosphorus loading to Lake Koochanusa was a fertilizer plant near Kimberley, British Columbia (Bonde and Bush, 1975). Daley and others (1980) cited this plant as a major source of orthophosphate and, to a lesser degree, ammonia nitrogen that has entered Kootenay Lake, which is about 230 km downstream from Libby Dam. The plant, in operation since 1953, discharged wastes to the St. Mary River, a tributary of the Kootenai River. Fertilizer production was doubled in 1962 and again increased in 1965. Water pollution control at the plant was improved in 1969, but it was not operating optimally until 1975 (Daley and others, 1980). The effect of the 1969 water-pollution control measures is evident in the history of orthophosphate loading in the Kootenai River at a station 6 km downstream from Libby Dam (streamflow-

sampling station 1, fig. 1). In 1968, 3,259 Mg of orthophosphate passed the station, but from 1969 to 1971, the annual loading of orthophosphate was less than 900 Mg (Bonde and Bush, 1975).




FIGURE 3.—Daily variation in surface altitude of Lake Koocanusa, 1972-80. \bar{X} denotes mean annual reservoir surface altitude, in meters.

TABLE 3.—Lake-filling and hydraulic-residence times of Lake Koocanusa, 1972-80

Year	Lake-filling time, in years				Hydraulic-residence time, in years			
	Annual	Monthly			Annual	Monthly		
		Mean ^a	Mini- mum	Maxi- mum		Mean ^a	Mini- mum	Maxi- mum
1972 ^b	0.14	0.17	0.04	0.52	0.14	0.14	0.02	0.37
1973	.22	.40	.10	.88	.33	.49	.11	1.29
1974	.28	.61	.09	1.28	.29	.33	.13	.67
1975	.37	.63	.11	1.13	.41	.78	.10	2.66
1976	.38	.71	.13	1.54	.38	.55	.13	1.56
1977	.64	.93	.26	1.64	.50	.59	.24	1.42
1978	.43	.75	.18	1.33	.48	.63	.24	1.28
1979	.66	1.08	.22	1.78	.62	.97	.22	2.08
1980	.52	.94	.17	1.47	.58	.78	.29	2.07

^aMean based on 12 monthly values, except for 1972.

^bApril through December only.

METHODOLOGY

Nutrient loadings that annually entered Lake Koocanusa were estimated to be the sum of loadings from gaged streamflow, ungaged drainage area, atmospheric deposition, and the Eureka sewage-treatment plant. Nutrient loadings from ground water and nutrient fluxes due to changes in reservoir bank storage were not quantified because of a lack of data.

The only nutrient loading estimated from gaged streamflow was for streamflow-sampling station 3 on the Kootenai River in British Columbia (fig. 1). The drainage area upstream from this station is 11,266 km² and the streamflow includes the effluent from the fertilizer plant at Kimberley. Annual nutrient loadings from this station were summations of daily loadings that are calculated as follows:

$$L = C \times Q \times f \quad (3)$$

where

L is daily loading, in megagrams;

C is daily nutrient concentration, in milligrams per liter;

Q is mean daily streamflow, in cubic meters per second; and

f is 0.08640, a factor to convert equation units to megagrams per day.

Daily nutrient concentrations were estimated from linear interpolation of nutrient concentrations actually measured in the gaged stream. Loadings of nutrients discharged from Lake Koocanusa were estimated with equation 3. The annual loadings of total phosphorus and total nitrogen that entered Lake Koocanusa prior to

impoundment were estimated by Bonde and Bush (1975) in a manner similar to equation 3.

Nutrient loading from the 12,005 km² of drainage area that was ungaged was estimated by the method of watershed-export coefficients of Reckhow, Beaulac, and Simpson (1980). Sufficient data were available from the Bull, Elk, and Tobacco Rivers to permit calculating loadings per unit area. Streamflow and water-quality data for 1972–77 were input to equation 3 and the results were summed to derive annual loadings of total phosphorus and total nitrogen. The annual loadings for each stream then were divided by their gaged drainage area. The median value of the derived watershed-export coefficients for the three streams was 0.144 kg·ha⁻¹·a⁻¹ (range = 0.022–0.796, number of samples = 13) for total phosphorus and 1.35 kg·ha⁻¹·a⁻¹ (range = 0.52–1.75, number of samples = 14) for total nitrogen. These two median watershed-export coefficients were multiplied by the ungaged drainage area of 1,200,502 ha to obtain annual loadings of total phosphorus and total nitrogen in kilograms. These values then were multiplied by 0.001 to convert them into megagrams.

Atmospheric deposition of total phosphorus and total nitrogen received by the surface of Lake Kocanusa was estimated using the product of the reservoir's mean annual surface area in hectares, a forest-atmospheric input coefficient in kilograms per hectare per year, and a 0.001 factor which yielded annual loading in megagrams. Forest-atmospheric input coefficients were 0.27 kg·ha⁻¹·a⁻¹ for total phosphorus and 0.99 kg·ha⁻¹·a⁻¹ for total nitrogen—the only values cited in Reckhow, Beaulac, and Simpson (1980) that were judged by this author as geographically applicable to the study area. Mean annual surface area, in hectares, was derived from statistical analysis of daily measurements of surface altitude. Mean annual reservoir surface altitude was converted to surface area using area-capacity curves for Lake Kocanusa.

Loadings of total phosphorus and total nitrogen from the sewage-treatment plant at Eureka, Montana (fig. 1), were obtained from the U.S. Environmental Protection Agency (1977). Annual loadings of 1.8 Mg of total phosphorus and 9.1 Mg of total nitrogen were measured during 1975 at the Eureka plant.

Variations in annual loadings of total phosphorus and total nitrogen were due, in part, to variations in annual streamflow. To remove this effect, each annual loading was divided by its respective annual streamflow to obtain a loading rate per unit volume of streamflow, expressed as megagrams per cubic kilometer of streamflow.

The relationship between influent loadings of total phosphorus and total nitrogen and the loadings of these

two nutrients discharged through Libby Dam was quantified with a nutrient-retention coefficient (Dillon and Rigler 1974), calculated thusly:

$$C = 1.0 - \frac{E}{I} \quad (4)$$

where

C is a nutrient-retention coefficient;

E is the mass of nutrient, in megagrams, discharged from the reservoir in a year; and

I is the mass of nutrient, in megagrams, that entered the reservoir in a year.

INFLUENT LOADINGS

Listed in tables 4 and 5 for total phosphorus and total nitrogen are the annual loadings and loading rates per unit volume of streamflow that entered Lake Kocanusa via gaged streamflow, ungaged drainage area, atmospheric deposition, and the Eureka sewage-treatment plant. From 1972 through 1975, the influent loading of total phosphorus ranged from 1,188 to 1,626 Mg. A large reduction occurred in 1976; the 1976 loading of total phosphorus was 39.4 percent of that estimated for 1975. For 1977–80, the loading of total phosphorus varied from 362 to 498 Mg. The loading rate per unit volume of streamflow for total phosphorus generally followed the same trend as the annual loadings of total phosphorus. The changes in total loading of total phosphorus primarily resulted from changes in loadings from

TABLE 4.—Annual loadings of total phosphorus that entered Lake Kocanusa, 1970–80^a

Year	Loading, in megagrams			Loading rate, in megagrams per cubic kilometer of streamflow
	Total ^b	Gaged inflow	Atmospheric	
1970	1,905	1,905	—	255.1
1971	1,924	1,924	—	162.1
1972	1,188	1,011	2.0	92.2
1973	1,626	1,449	2.1	194.1
1974	1,485	1,307	3.4	106.8
1975	1,304	1,126	3.4	129.2
1976	514	336	3.7	44.0
1977	362	184	3.4	58.3
1978	498	320	3.7	49.0
1979	416	238	3.8	59.8
1980	428	250	4.0	45.2

^aPreimpoundment loadings for 1970–71 calculated for streamflow-sampling station 1, located 6 kilometers downstream from Libby Dam.

^bTotal loadings for 1972–80 include constant annual loadings of 172.6 megagrams from ungaged inflow and 1.8 megagrams from Eureka sewage-treatment plant.

TABLE 5.—Annual loadings of total nitrogen that entered Lake Kooconusa, 1970-80^a

Year	Loading, in megagrams			Loading rate, in megagrams per cubic kilometer of streamflow
	Total ^b	Gaged inflow	Atmospheric	
1970	2,825	2,825	--	378.3
1971	4,057	4,057	--	341.8
1972	4,679	3,042	7.5	363.5
1973	3,228	1,590	7.8	385.3
1974	4,051	2,409	12.4	291.3
1975	2,798	1,156	12.3	277.2
1976	3,089	1,445	13.5	264.4
1977	2,451	809	12.6	395.0
1978	2,891	1,248	13.4	284.4
1979	2,707	1,063	14.0	389.5
1980	3,258	1,613	14.6	343.6

^aPreimpoundment loadings for 1970-71 calculated for streamflow-sampling station 1, located 6 kilometers downstream from Libby Dam.

^bTotal loadings for 1972-80 include constant annual loadings of 1,621 megagrams from ungaged inflow and 9.1 megagrams from Eureka sewage-treatment plant.

gaged streamflow, which reflected the reduced loadings of total phosphorus discharged by the fertilizer plant. Data on loadings of total phosphorus and total nitrogen during 1973-78 in the St. Mary River upstream and downstream from the fertilizer plant (streamflow-sampling stations 4 and 5, fig. 1) readily showed the reduced loading of total phosphorus that commenced in 1976 (table 6). The loadings from the Eureka sewage-treatment plant and atmospheric deposition were too small to significantly alter the total loading. Ungaged drainage areas contributed a constant yearly load of 172.6 Mg of total phosphorus. Variations in loadings from ungaged drainages may have significantly affected the total loading; however, such variations could not be estimated because of the lack of data.

The declining trend in influent loading that was readily apparent for total phosphorus did not occur for loadings of total nitrogen. During 1972-80, influent loadings of total nitrogen ranged from 2,451 to 4,679 Mg; nitrogen loading was not largely reduced in 1976 (table 5) as was total phosphorus loading. Variations in the loading of total nitrogen were primarily due to variations in loading of total nitrogen in gaged inflow, because ungaged inflow was estimated as a constant loading and loadings from the Eureka sewage-treatment plant and atmospheric deposition were too small to significantly affect total loading. The loading rate per unit volume of streamflow for total nitrogen showed no definite trend (table 5). Water-pollution-control measures instituted at the fertilizer plant apparently had little effect

TABLE 6.—Comparison of 1973-78 loadings of total phosphorus (TP) and total nitrogen (TN) at two stations on the St. Mary River, one upstream and one downstream from the fertilizer plant at Kimberley, British Columbia^a

Year	Annual loading, in megagrams				Percentage upstream loading is of downstream loading	
	Upstream station ^b		Downstream station ^c		TP	TN
	TP	TN	TP	TN		
1973	8.1	175.4	1,806	532.5	22,270	303.7
1974	15.5	264.1	1,671	937.3	10,770	354.9
1975	7.9	196.1	1,215	579.4	15,340	295.4
1976	10.5	259.0	459.4	433.4	4,388	167.3
1977	3.6	125.0	165.2	291.3	4,538	233.0
1978	8.9	178.1	226.5	411.2	2,548	230.9

^a Loading data from James Helms (U.S. Army Corps of Engineers, written commun., 1981).

^b Streamflow-sampling station 4 (fig. 1)

^c Streamflow-sampling station 5 (fig. 1)

on the amount of total nitrogen carried per unit volume of streamflow, because loadings of total nitrogen upstream and downstream from the fertilizer plant did not substantially change during 1973-78 (table 6).

LOADINGS DISCHARGED FROM LAKE KOOCANUSA

After closure of Libby Dam in March 1972, loadings of total phosphorus substantially declined downstream from the dam; reductions in loading of total nitrogen were not as apparent, although some reduction occurred (table 7). Loading rates per unit volume of streamflow for total phosphorus and total nitrogen showed trends similar to those for their annual loadings. The variations in loading of total phosphorus and total nitrogen discharged from Lake Kooconusa were due, in part, to the water-pollution-control measures instituted at the fertilizer plant, which were previously discussed. However, the loadings that were input at the upstream end of the reservoir were subjected to sedimentation and chemical-biological transformations prior to their release from Libby Dam. Because of these two processes, variations in loads discharged from the dam cannot be directly attributed to variations in loads from the fertilizer plant.

NUTRIENT RETENTION IN LAKE KOOCANUSA

The combined effects within the reservoir on influent loadings of total phosphorus and total nitrogen were estimated with nutrient-retention coefficients (table 8).

TABLE 7.—Annual loadings of total phosphorus (TP) and total nitrogen (TN) discharged from Lake Kooconusa, 1970–80^a

Year	Loading, in megagrams		Loading rate, in megagrams per cubic kilometer of streamflow	
	TP	TN	TP	TN
1970	1,905	2,825	255.1	378.3
1971	1,924	4,057	162.1	341.8
1972	997	4,004	78.5	315.1
1973	554	1,702	98.5	302.5
1974	706	3,378	53.2	254.8
1975	326	1,876	35.9	206.3
1976	359	2,529	30.8	216.5
1977	125	2,359	15.9	299.4
1978	82	2,229	9.0	246.7
1979	45	2,201	6.1	295.3
1980	50	1,626	5.9	191.5

^aPreimpoundment loadings for 1970–71 calculated for streamflow-sampling station 1, located 6 kilometers downstream from Libby Dam; loadings for 1972–80 calculated for streamflow-sampling station 2, located 1.1 kilometers downstream from Libby Dam.

These coefficients indicate that total phosphorus was more readily retained in the reservoir than was total nitrogen. During 1972–80, an average of 63 per cent of influent total phosphorus was retained, whereas 25 percent of the total nitrogen loading was retained.

The large retention of total phosphorus in Lake Kooconusa was partly accounted for by the tendency for phosphorus to adsorb to clay minerals (Golterman 1975; Lee, 1970). Upstream from Lake Kooconusa, the Kootenai River traverses sedimentary and glacial deposits containing clay and silt (Crozier and Leinweber, 1975). Eroded sediments were transported into the reservoir and were deposited; part of the influent load of total phosphorus was, thereby, deposited. The efficiency of sediment trapping by reservoirs was empirically related to lake-filling time by Brune (1953). Based on Brune's graphs and Lake Kooconusa's theoretical lake-filling time of 0.67 year, the sediment trapping efficiency of the reservoir exceeds 95 percent. This value was verified by suspended-sediment measurements at streamflow-sampling stations 1 and 2, located downstream from Libby Dam. The average annual loading of suspended sediment per cubic kilometer of streamflow was 131,792 Mg from 1968 through 1971 and 9,288 Mg from 1972 through 1975. Postimpoundment loadings of suspended sediment, therefore, were reduced an average of 122,504 Mg, or by 93 percent, following impoundment of Lake Kooconusa.

The chemistry of phosphorus in Lake Kooconusa sediments was investigated by Iskandar and Shukla (1981). Their samples indicated a very small organic matter content, which they stated was characteristic of oligo-

TABLE 8.—Nutrient-retention coefficients for total phosphorus and total nitrogen in Lake Kooconusa, 1972–80

Year	Nutrient-retention coefficient	
	Total phosphorus	Total nitrogen
1972	0.16	0.14
1973	.66	.47
1974	.52	.17
1975	.75	.33
1976	.30	.18
1977	.66	.04
1978	.84	.23
1979	.89	.19
1980	.88	.50

trophic lake sediments. The total phosphorus content of the silty clay sediments was composed of 87 to 98 percent inorganic phosphorus. These authors determined experimentally that the sediments had limited ability to adsorb additional phosphorus and the sediments desorbed only small amounts of phosphorus. Iskandar and Shukla concluded that Lake Kooconusa sediments function as a phosphorus sink. This conclusion appears reasonable because of the 7,821 Mg of total phosphorus that entered the reservoir during 1972–80, the reservoir discharged 3,244 Mg during the same period.

Part of the loading of total nitrogen also may have sorbed to clay minerals that were deposited in the reservoir. Although nitrate has little tendency to sorb to clay minerals (Lee, 1970), Keeney (1972) reported that organic nitrogen compounds readily sorb to clays and other inorganic colloids. During 1972–80, total nitrogen loading can be divided, on the average, into the following nitrogen forms: Nitrite plus nitrate, 45.6 percent; organic nitrogen, 43.1 percent; and ammonia, 8.2 percent. Presumably, less than 43.1 percent of the total nitrogen loading would have been adsorbed and deposited during 1972–80.

Further consideration of the retention of nitrogen and phosphorus loadings in Lake Kooconusa was difficult because both are nonconservative nutrients required by algae for photosynthesis and growth. The two nutrients are thereby involved in a complex sequence of uptake, utilization, and remineralization. For example, a blue-green alga, *Aphanizomenon flos-aquae*, found in Lake Kooconusa has been reported by Fogg (1974) to fix atmospheric nitrogen and, thereby, may have provided an additional source of nitrogen to the reservoir. However, an undetermined amount of nitrogen was removed to the reservoir's sediment via denitrification reactions. Although the chemically and biologically induced processes that affect nitrogen and phosphorus within aqua-

tic ecosystems are widely recognized, they are infrequently quantified. Detailed nutrient budgets that quantify these processes require data that were not available for Lake Kooconusa.

PRIMARY PRODUCTIVITY

METHODOLOGY

Primary productivity was measured at the four limnological stations (fig. 1) commencing in late July 1972; sampling was discontinued at Pinkham after July 1976. Most of the 313 samples of primary productivity were obtained during May through October; only nine were taken in April and nine in November. Pinkham and Boundary generally were not sampled until late May or early June, because reservoir drawdown created insufficient depth for sampling prior to this time.

Primary productivity was measured with the carbon-14 method for phytoplankton described by Greeson, Ehlike, Irwin, Lium, and Slack (1977). Five water samples were retrieved from depths at which the light intensity was equal to or larger than 0.1 percent of that incident at the reservoir surface. Light measurements were made with an unfiltered submersible photometer immediately prior to retrieval of the water samples. The water samples were then transferred to light and dark BOD bottles and inoculated with radioactive carbonate ($^{14}\text{CO}_3^{2-}$). They were then incubated for 3 to 4 hours at the depth from which they were withdrawn. The light intensities used for incubation generally included five of the following seven values: 90, 60, 30, 15, 5, 1, and 0.1 percent. During the study, incubations generally were conducted during the period spanning 0900 to 1500 hours.

The value calculated for each water sample represents daily primary productivity, expressed as milligrams of carbon fixed per cubic meter ($\text{mg}\cdot\text{C}\cdot\text{m}^{-3}$). Values of daily primary productivity measured within the sampled depth profile were integrated to obtain daily areal primary productivity, expressed as milligrams of carbon fixed per square meter ($\text{mg}\cdot\text{C}\cdot\text{m}^{-2}$).

Only primary productivity by phytoplankton was measured in this study. No estimates of periphyton primary productivity were available for Lake Kooconusa. However, in large and deep lakes one generally can assume that the majority of annual primary productivity is by phytoplankton (Likens, 1975).

Annual areal primary productivity, in grams of carbon fixed per square meter ($\text{g}\cdot\text{C}\cdot\text{m}^{-2}$), was estimated for Lake Kooconusa to permit comparisons with other lakes and reservoirs. Annual areal primary productivity was calculated by integrating the area under a curve defined by values of total daily primary productivity for the reservoir. The total daily values were derived

with the following equation :

$$A = (R_F \times P_F) + (R_T \times P_T) + (R_P \times P_P) + (R_B \times P_B) \quad (5)$$

where

A is total daily primary productivity, in grams of carbon fixed per square meter;

R_F is daily primary productivity, in grams of carbon fixed per square meter, at Forebay station (fig. 1);

R_T is daily primary productivity, in grams of carbon fixed per square meter, at Tenmile station;

R_P is daily primary productivity, in grams of carbon fixed per square meter, at Pinkham station;

R_B is daily primary productivity, in grams of carbon fixed per square meter, at Boundary station;

P_F is fraction of reservoir surface area associated with Forebay primary productivity value;

P_T is fraction of reservoir surface area associated with Tenmile primary productivity value;

P_P is fraction of reservoir surface area associated with Pinkham primary productivity value; and

P_B is fraction of reservoir surface area associated with Boundary primary productivity value.

Days on which primary productivity was measured were used in equation 5; however, samples were not taken during much of the winter so this period required estimation. Based on primary productivity measurements at Forebay in November and December, the following estimates of daily areal primary productivity, in grams of carbon fixed per square meter, were used for each limnological station during winter months: November, December, and March, 0.01; January and February, 0.005. The months of April through July occasionally lacked a measured value of daily areal primary productivity at one or more limnological stations. In such instances, the average daily areal primary productivity at the limnological station during the missing month was used as input to equation 5.

SEASONAL TREND

Daily areal primary productivity varied considerably among limnological stations and years and ranged from 0.4 to 420.0 $\text{mg}\cdot\text{C}\cdot\text{m}^{-2}$ (table 9). Statistical tests used to detect significant differences among limnological stations and years were not performed because of differences in the period sampled at each limnological station. Instead, the mean, minimum, and maximum values of daily areal primary productivity for each month are given (table 9). With the four limnological stations combined, August had the largest mean daily areal primary productivity and November had the smallest.

Mean values of daily areal primary productivity for April through June were substantially smaller at

PRIMARY PRODUCTIVITY

TABLE 9.—*Statistical summary of daily areal primary productivity and euphotic zone depth at the four limnological stations, Lake Kooconusa, 1972-80*

Station	Month	Daily areal primary productivity, in milligrams of carbon fixed per square meter			Euphotic zone depth, in meters			Number of Samples
		Mean	Minimum	Maximum	Mean	Minimum	Maximum	
Boundary	June	82.6	0.4	150.0	4.9	0.4	11.4	9
	July	135.1	75.0	340.0	7.3	2.1	12.2	17
	Aug.	179.7	96.0	290.0	9.6	5.8	13.5	15
	Sept.	136.4	20.0	230.0	10.2	5.5	16.0	17
	Oct.	90.3	24.0	220.0	8.7	5.9	14.8	15
Pinkham	Apr.	75.0	75.0	75.0	1.4	1.4	1.4	1
	May	10.2	.5	36.0	1.1	.4	1.8	6
	June	83.0	18.0	230.0	2.8	1.1	4.9	5
	July	119.7	28.0	270.0	5.1	1.7	7.8	7
	Aug.	131.0	86.0	170.0	8.4	5.4	10.7	6
	Sept.	154.3	96.0	250.0	9.7	8.5	10.5	6
	Oct.	65.7	24.0	120.0	7.6	5.8	10.3	7
Tenmile	Nov.	6.7	1.4	12.0	4.8	.5	9.2	2
	Apr.	119.8	57.0	250.0	2.6	1.4	3.7	4
	May	109.4	16.0	390.0	4.6	.6	9.5	14
	June	171.3	100.0	280.0	5.8	1.7	10.7	15
	July	147.7	41.0	220.0	8.0	2.3	14.5	15
	Aug.	152.2	43.0	220.0	11.3	5.9	18.0	16
	Sept.	138.2	20.0	230.0	12.5	8.5	18.4	17
Forebay	Oct.	109.1	44.0	200.0	11.7	7.7	19.8	16
	Nov.	20.0	5.1	43.0	7.2	4.9	11.2	3
	Apr.	168.5	23.0	340.0	4.0	2.7	4.7	4
	May	139.4	62.0	320.0	6.0	2.3	11.2	14
	June	165.1	92.0	400.0	6.3	2.3	10.8	15
	July	141.9	12.0	370.0	7.8	1.8	12.2	16
	Aug.	171.2	63.0	420.0	11.4	7.1	18.0	16
All	Sept.	126.3	37.0	190.0	11.8	9.0	16.0	15
	Oct.	107.3	40.0	230.0	11.0	7.6	16.8	15
	Nov.	31.0	3.2	69.0	5.2	3.7	7.9	4
	Apr.	136.4	23.0	340.0	3.1	1.4	4.7	9
	May	104.2	.5	390.0	4.5	.4	11.2	34
	June	141.0	.4	400.0	5.5	.4	11.4	44
	July	138.6	12.0	370.0	7.4	1.7	14.5	55
All	Aug.	163.4	43.0	420.0	10.5	5.4	18.0	53
	Sept.	136.2	20.0	250.0	11.3	5.5	18.4	55
	Oct.	97.7	24.0	230.0	10.1	5.8	19.8	54
	Nov.	22.0	1.4	69.0	5.8	.5	11.2	9
	Apr.-Nov.	128.5	.4	420.0	8.3	.4	19.8	313

Boundary and Pinkham than at Tenmile and Forebay. This condition is attributed to the fact that areal primary productivity is dependent on euphotic zone depth and the depths of the euphotic zone during April through June were smallest at Boundary and Pinkham. Mean euphotic zone depths during April and May at Pinkham were equal to or less than 1.4 m, whereas they were equal to or greater than 2.6 m at Tenmile and Forebay. Although not measured in April and May, euphotic zone depths at Boundary probably were similar to those at Pinkham. The shallow euphotic zones during April and May occurred when the reservoir received large loadings of suspended sediment that were reported by Bonde and Bush (1975) to be historically carried by the Kootenai River during spring runoff.

The monthly based means for daily areal primary productivity in table 9 failed to show the variation among years; such variation is evident in table 10. Because sampled periods did not coincide, statistical tests were not used to locate significant differences among the years. Mean daily areal primary productivity varied from 106.7 to 165.5 $\text{mg}\cdot\text{C}\cdot\text{m}^{-2}$, but the overall range during the 9 years was 0.4 to 420.0 $\text{mg}\cdot\text{C}\cdot\text{m}^{-2}$. The variation in daily areal primary productivity within each year at the four limnological stations is apparent from figure 4. No common pattern among stations or years could be discerned, owing to the wide variation in daily areal primary productivity. Such variation was due to the effect of environmental variables on daily areal primary productivity; however, evaluation of the effects of such variables on daily areal primary productivity was beyond the scope of this report.

ANNUAL TREND

Annual areal primary productivity ranged from 23.2 to 38.5 and averaged 29.5 $\text{g}\cdot\text{C}\cdot\text{m}^{-2}$ during 1972-80 (table 11). The 95-percent confidence interval for the mean was determined to be 25.6 to 33.4 $\text{g}\cdot\text{C}\cdot\text{m}^{-2}$. The years with annual primary productivity outside of the confidence interval included 1973-76 and 1980.

Primary productivity in Lake Kooconusa can be compared to general ranges of phytoplankton primary productivity of lakes in various trophic categories that were listed in Wetzel (1975). Wetzel listed mean primary productivity, in milligrams of carbon fixed per square meter per day, which is identical to average daily areal primary productivity listed in table 11. Three of the trophic categories listed by Wetzel (1975), based on mean primary productivity, in milligrams of carbon fixed per square meter per day, were: Oligotrophic, 50-300; mesotrophic, 250-1,000; and eutrophic, greater than 1,000. Lake Kooconusa was clearly in the oligotrophic category, based on its average daily areal primary productivity that ranged from 63.6 to 105.5 $\text{mg}\cdot\text{C}\cdot\text{m}^{-2}$ (table 11).

TABLE 10.—Statistical summary of daily areal primary productivity in Lake Kooconusa, 1972-80

Year	Period sampled	Daily areal primary productivity, in milligrams of carbon fixed per square meter			Number of samples
		Mean	Minimum	Maximum	
1972	July 24 - Nov. 15	122.8	1.4	290.0	19
1973	Apr. 24 - Nov. 30	120.5	.5	390.0	28
1974	Apr. 23 - Nov. 20	106.7	2.1	400.0	51
1975	Apr. 16 - Oct. 30	115.5	.4	320.0	48
1976	Apr. 12 - Oct. 28	114.1	23.0	420.0	39
1977	May 4 - Oct. 26	139.8	20.0	280.0	30
1978	May 5 - Oct. 25	146.2	80.0	230.0	33
1979	May 15 - Oct. 17	143.6	62.0	370.0	32
1980	May 14 - Oct. 16	165.5	45.0	340.0	33

RELATIONSHIP OF ANNUAL AREAL PRIMARY PRODUCTIVITY AND SELECTED ENVIRONMENTAL VARIABLES

Variations in annual areal primary productivity are caused by numerous environmental variables that directly or indirectly control the primary productivity of phytoplankton. The relationship of environmental variables and annual areal primary productivity in Lake Kooconusa was investigated with correlation and multiple regression analyses. Annual-based values for 12 environmental variables (table 12) were derived from the data base for use in these statistical analyses. Although table 12 is not an exhaustive listing, each of the 12 variables has a direct or indirect causal relationship with phytoplankton primary productivity.

Correlation coefficients for each environmental vari-

TABLE 11.—Annual areal primary productivity in Lake Kooconusa, 1972-80

Year	Annual areal primary productivity, in grams of carbon fixed per square meter	Average daily areal primary productivity, in milligrams of carbon fixed per square meter
1972 ^a	27.1	74.2
1973	38.5	105.5
1974	25.3	69.3
1975	24.4	66.8
1976	23.2	63.6
1977	30.6	83.8
1978	30.7	84.1
1979	30.0	82.2
1980	35.3	96.7
1972-80	29.5	80.7

^aIncludes March 21 to December 31.




FIGURE 4.—Daily values of areal primary productivity at the four limnological stations, Lake Kooconasa, 1972–80.

able and annual areal primary productivity are listed in table 13. The two largest correlation coefficients exceeded 0.500 and were for annual nitrogen retention coefficient (X12) and mean annual euphotic zone depth (X1). However, none of the correlation coefficients were significantly different from 0.00 when judged against a significance level of 0.05 or less. Therefore, no single

environmental variable was usable as a predictor of annual areal primary productivity. Combinations of the 12 environmental variables, therefore, were analyzed by multiple linear regression analysis, using annual areal primary productivity as the dependent variable.

Variable combinations were initially screened with the RSQUARE procedure of the Statistical Analysis

TABLE 12.—Statistical summary of environmental variables used in correlation and multiple regression analyses of annual areal primary productivity in Lake Kooconusa, 1972-80

Variable no.	Variable label	Mean	Minimum	Maximum
Y	Annual areal primary productivity, in grams carbon fixed per square meter	29.5	23.2	38.5
X1	Mean annual euphotic zone depth, in meters	8.5	5.9	12.5
X2	Maximum annual euphotic zone depth, in meters	13.8	10.7	19.8
X3	Mean annual water temperature at 3 meters depth, in degrees Celsius	14.6	12.2	16.7
X4	Maximum annual water temperature at 3 meters depth, in degrees Celsius	20.5	18.8	22.0
X5	Mean monthly lake filling time, in years	.69	.17	1.08
X6	Range of monthly lake filling time, in years	1.14	.48	1.56
X7	Mean monthly hydraulic residence time, in years	.58	.14	.97
X8	Range of monthly hydraulic residence time, in years	1.33	.36	2.57
X9	Annual areal phosphorus loading, in grams per square meter	8.3	2.8	20.7
X10	Annual areal nitrogen loading, in grams per square meter	29.1	19.1	61.7
X11	Annual phosphorus retention coefficient	.63	.16	.89
X12	Annual nitrogen retention coefficient	.25	.04	.50

System (SAS Institute, Inc., 1979). This computer procedure determines all possible linear regressions for a dependent variable and a set of independent variables, and then lists the coefficient of multiple correlation for each regression. Although 12 independent variables were available, there were only nine observations on each variable, which required that less than nine variables be used in each regression. Based on the advice of R. K. Steinhurst (University of Idaho, oral commun., 1981), only two- and three-variable regressions were developed from the data base. Those two- and three-variable regressions with the largest coefficient of multiple determination were selected for further analysis using the GLM procedure in the Statistical Analysis System. This computer procedure provides the parameter estimates for a regression model, the analysis of variance table, hypothesis tests, and residuals.

The two- and three-variable regression models that were derived from the 12 environment variables and that had the largest coefficient of multiple determination are listed in table 14. In the two-variable model, annual areal primary productivity varied directly with mean annual euphotic zone depth and annual areal phosphorus loading. In the three-variable model, the dependent variable varied directly with mean annual euphotic zone depth and annual nitrogen retention coefficient and it varied inversely with maximum annual euphotic zone

TABLE 13.—Correlation coefficients between annual areal primary productivity and environmental variables X1 through X12^a

[Number of samples is nine]

Variable no.	Correlation coefficient	Significance level of correlation coefficient
X1	0.522	0.150
X2	.355	.349
X3	.392	.296
X4	.418	.263
X5	.100	.800
X6	-.106	.786
X7	.187	.630
X8	-.012	.976
X9	.185	.633
X10	.043	.913
X11	.493	.178
X12	.605	.084

^aVariables defined in table 12.

depth. Mean annual euphotic zone depth was the dominant independent variable in both regression models, based on the relative magnitude of standardized partial regression coefficients. The two regression models account for 62.0 and 78.4 percent of the variation in the dependent variable. The overall regression models and their partial regression coefficients were significantly different from 0.00 at significance levels less than 0.061.

The residuals for each regression model were graphically analyzed according to methods of Draper and Smith (1966). These analyses revealed no apparent violations in the assumptions of regression analysis. Scatter plots of each independent variable and the dependent variable exhibited linear trends; therefore, the independent variables were not transformed.

The two regression models in table 14 provided an empirical method for predicting annual areal primary productivity in Lake Kooconusa. The models were developed with a statistical technique that only quantifies the variation between dependent and independent variables. The method does not prove the existence of cause-and-effect relationships between the variables. The relationships may be spurious if the variables varied systematically, but independent of one another. Therefore, the two regression models were examined further to determine if cause-and-effect relationships were plausible between the dependent and independent variables. Correlations between the dependent and independent variables were examined with scatterplots (fig. 5) and the temporal variation of the variables was plotted (fig. 6) to assess systematic covariation.

Mean annual euphotic zone depth, the dominant variable in both regression models, was directly related to

TABLE 14.—Multiple regression models for prediction of annual areal primary productivity in Lake Kooconusa

TWO INDEPENDENT VARIABLES			
Model			
$Y = 7.725 + (2.020 \times X1) + (0.548 \times X9)$			
where			
Y is annual areal primary productivity, X1 is mean annual euphotic zone depth, and X9 is annual areal phosphorus loading.			
Multiple correlation coefficient = 0.788 Coefficient of multiple determination = 0.620 Significance level of overall regression = 0.055			
Vari- able	Partial regression coefficient	Standardized partial regression coefficient	Significance level of partial regression coefficient
X1	2.020	0.934	0.023
X9	0.548	0.720	0.057
THREE INDEPENDENT VARIABLES			
Model			
$Y = 23.852 + (4.310 \times X1) + (-2.583 \times X2) + (18.120 \times X12)$			
where			
Y is annual areal primary productivity, X1 is mean annual euphotic zone depth, X2 is maximum annual euphotic zone depth, and X12 is annual nitrogen retention coefficient.			
Multiple correlation coefficient = 0.886 Coefficient of multiple determination = 0.784 Significance level of overall regression = 0.041			
Vari- able	Partial regression coefficient	Standardized partial regression coefficient	Significance level of partial regression coefficient
X1	4.310	1.993	0.034
X2	-2.583	-1.662	0.060
X12	18.120	0.552	0.049

annual areal primary productivity (table 14). This relationship appeared plausible because areal primary productivity might be expected to increase if the depth of the euphotic zone increased. Mean annual euphotic zone depth plotted a positive linear trend with annual areal primary productivity, although the value for 1973 appeared to be outside of the trend (fig. 5). Over time, mean annual euphotic zone depth had an increasing trend similar to that of annual areal primary productivity, excepting the large 1973 value of the latter variable (fig. 6).

Maximum annual euphotic zone depth was inversely related to annual areal primary productivity in the three-variable regression model (table 14). This result does not seem plausible because mean and maximum euphotic zone depths were positively correlated (correlation coefficient = 0.953, significance level less than 0.0001, number of samples = 9) and, therefore, would be expected to affect annual areal primary productivity in a similar manner. Maximum annual euphotic zone depth plotted a positive linear trend with annual areal primary productivity, although the values for 1973 and 1979 appeared to plot outside of the trend (fig. 5). Similar temporal trends, excepting 1973, were exhibited by

maximum annual euphotic zone depth and annual areal primary productivity (fig. 6). Based on this conflicting evidence, the occurrence of this variable in the regression model appeared to be spurious.

Annual areal phosphorus loading was directly related to annual areal primary productivity in the two-variable regression model (table 14). The relationship appeared to be plausible because increased phosphorus loading might be expected to result in increased annual areal primary productivity. The scatterplot of annual areal phosphorus loading showed a positive linear trend, which was due largely to the location of the 1973 value (fig. 5). If the 1973 value were deleted, the trend would likely have been negatively linear. Annual areal phosphorus loading had a decreasing trend with time (fig. 6), as would be expected from the reductions in total phosphorus loading listed in table 4. During 1972–76 and 1977–79, annual areal phosphorus loading and annual areal primary productivity covaried systematically, which largely explains the direct relationship derived by the regression modeling technique.

In the three-variable regression model, annual nitrogen retention coefficient was directly related to annual areal primary productivity (table 14). This relationship appeared to be intuitively correct, because additional retention of nitrogen might be expected to increase annual areal primary productivity. However, the positive linear trend in the scatterplot (fig. 5) was produced by the values for 1973 and 1980; if these two were not present, the trend would have been negatively linear. Figure 6 shows that the annual nitrogen retention coefficient varied sporadically with time, but during 1972–74 and 1979–80 it varied systematically with annual areal primary productivity. The inclusion of annual nitrogen retention coefficient into the regression model appeared to be largely dependent upon the bias created by its two largest values, which corresponded to the two largest values of annual areal primary productivity.

Of the four independent variables, only mean annual euphotic zone depth appeared likely to have a cause-and-effect relationship with annual areal primary productivity. The direct relationships of annual areal phosphorus loading and annual nitrogen retention coefficient with annual areal primary productivity were strongly biased by the coincidence of their largest values. The inverse relationship of maximum annual euphotic zone depth and annual areal primary productivity was deemed spurious because it was intuitively incorrect.

Mean annual euphotic zone depth was indicative of the depth of light penetration into Lake Kooconusa. The attenuation of light by water can be quantified with an extinction coefficient that composites effects due to absorption of light by water and the scattering of light by suspended and dissolved matter. The extinction coef-




FIGURE 5.—Scatterplots of annual areal primary productivity versus maximum and mean annual euphotic zone depths, annual areal phosphorus loading, and annual areal nitrogen retention coefficient for Lake Kooconusa, 1972-80.




FIGURE 6.—Annual variation of mean and maximum euphotic zone depths, areal primary productivity, areal phosphorus loading, and nitrogen retention coefficient for Lake Kooconusa, 1972–80.

ficient within the euphotic zone of Lake Kooconusa and the surface illumination were independent variables in a multiple regression derived by Woods (1979) to predict daily areal primary productivity measured in Lake Kooconusa from 1972 to 1975. Woods used 93 daily values instead of the 9 annual values reported here. The regression model is listed in table 15; derivation

of the variables was discussed by Woods (1979). The regression model showed that daily areal primary productivity was inversely related to euphotic extinction coefficient and directly related to surface illumination. These relationships appeared plausible because a reduction in euphotic zone depth caused by an increase in euphotic extinction coefficient would likely reduce daily

TABLE 15.—Multiple regression model developed by Woods (1979) for prediction of daily areal primary productivity in Lake Kooconusa

Model			
$Y = -62.12 + (-12.95 \times X1) + (0.17 \times X2) + (1.04 \times X3) + (2.12 \times X4) + (0.01 \times X5) + (0.05 \times X6)$			
where			
Y is daily areal primary productivity, in milligrams carbon fixed per square meter;			
X1 is euphotic extinction coefficient;			
X2 is areal loading, in meters per year;			
X3 is euphotic dissolved-solids concentration, in milligrams per liter;			
X4 is stability of primary thermocline, in grams per square centimeter;			
X5 is surface illumination, in foot-candles ^a ; and,			
X6 is hydraulic-residence time, in days.			
Multiple correlation coefficient = 0.707			
Coefficient of multiple determination = 0.500			
Significance level of overall regression = <0.001			
Vari- able	Partial regression coefficient	Standardized partial regression coefficient	Significance level of partial regression coefficient
X1	-12.95	-0.677	<0.001
X2	0.17	0.391	<0.001
X3	1.04	0.315	<0.001
X4	2.12	0.276	<0.005
X5	0.01	0.265	<0.005
X6	0.05	0.208	<0.025

^aTo convert foot-candle to lux, multiply by a factor of 10.76.

areal primary productivity; increased surface illumination should increase daily areal primary productivity. Results of multiple regression analyses for daily and annual values were consistent in that variables related to light penetration into the euphotic zone were important predictors of areal primary productivity.

Woods (1979) concluded from analyses of data collected from 1972-75 at Lake Kooconusa that the quantity of light received by phytoplankton was the major environmental influence on primary productivity in the reservoir—although no single environmental variable accounted for all variation. Three processes were identified as having had an effect on the light available to phytoplankton. First, a weak thermal structure in the reservoir allowed phytoplankton to be circulated out of the euphotic zone into light conditions inadequate for photosynthesis to exceed respiration. The operational schedule of the reservoir, in conjunction with large seasonal inflow and outflow, produced the weak thermal structure. Second, the largely turbid inflows during spring runoff substantially reduced euphotic zone depths and, thereby, intensified the problem of light limitation experienced by phytoplankton that was circulated out of the euphotic zone. Third, the phytoplanktonic light environment was largely dependent upon seasonally and meteorologically induced variations in incident solar radiation.

The foregoing conclusions of Woods (1979) also may be applicable to the environmental control of daily areal primary productivity during 1976-80 in Lake Kooconusa. However, the data analyses necessary to determine the environmental control of daily areal primary productivity during 1976-80 were beyond the

scope of this report. Evaluation of the relationship of daily values of primary productivity in conjunction with concurrent values for physical, chemical, and biological variables likely would yield valuable insight into the reasons why Lake Kooconusa was oligotrophic during 1972-80, in spite of its potentially eutrophic loadings of nitrogen and phosphorus.

CHLOROPHYLL *a* DISTRIBUTION WITHIN THE WATER COLUMN

Based on his analysis of data collected during 1972-75 at Lake Kooconusa, Woods (1979) hypothesized that phytoplankton was circulated beneath the euphotic zone. To test this hypothesis the limnological sampling program at Lake Kooconusa was modified in the 1980 water year. In that chlorophyll *a* is a primary photosynthetic pigment found in all algae (Wetzel, 1975), its distribution in the water column was used to test this hypothesis. Such tests were not feasible prior to 1980 because chlorophyll *a* was not sampled beneath the euphotic zone prior to 1980. In 1980, however, chlorophyll *a* was sampled at two points beneath the euphotic zone—one each at two and three times the depth of 0.1 percent light. In addition, five samples of chlorophyll *a* were collected in the euphotic zone—one each at the carbon-14 incubation depths. Prior to mid-May 1978 chlorophyll *a* in Lake Kooconusa was analyzed according to methods of Slack, Averett, Greeson, and Lipscomb (1973). This procedure did not correct for the degradation products of chlorophyll *a*; hence, one could not ascertain if the results were representative of living or dead chlorophyll-bearing organisms. Following mid-May 1978, chlorophyll *a* sampled from Lake Kooconusa was determined according to methods of Greeson, Ehlike, Irwin, Lium, and Slack (1977). These results indicated the amount of chlorophyll *a* that was contained in living chlorophyll-bearing organisms.

Chlorophyll *a* concentrations were sampled at Forebay and Tenmile in 1980. The volume of chlorophyll *a* in each of three depth strata then was determined from a plot of chlorophyll *a* concentration versus depth (fig. 7). The percentage of the total volume of chlorophyll *a* in the sampled water column then was determined for each of the three depth strata. The three depth strata were as follows: Surface to 1.0 percent light depth, 1.0 percent to 0.1 percent light depth, and 0.1 percent light depth to a depth equal to the depth of 0.1 percent light plus one-half of the depth difference between 1.0 percent and 0.1 percent light depth. The 1.0 percent light depth was chosen as the lower boundary of the upper stratum because the 1.0 percent light depth is often defined as the lower boundary of the euphotic zone (Greeson and others, 1977).

The 0.1 percent light depth was chosen as a boundary because it was often the depth at which the lowermost carbon-14 sample was incubated in Lake Kooacanusa.

The percentage of chlorophyll *a* in each stratum varied during the sampled interval; the 95-percent confi-

dence intervals indicate that the middle stratum varied less than the other two strata (table 16). The mean percentage distribution of chlorophyll *a* in the three strata was similar at both limnological stations. On the average, less than one-half of the sampled chlorophyll




FIGURE 7.—Typical diagram used to quantify the percentage distribution of chlorophyll *a* within the water column of Lake Kooacanusa.

TABLE 16.—Percentage distribution of chlorophyll *a* in three depth strata at the Forebay and Tenmile stations in 1980

Station	Date sampled	Upper stratum ¹	Middle stratum ²	Lower stratum ³	
Forebay	May 15	24.5	13.0	62.5	
	May 30	26.5	18.0	55.5	
	June 11	38.1	16.4	45.5	
	June 30	34.1	13.7	52.2	
	July 10	41.4	21.1	37.5	
	July 24	63.9	20.5	15.6	
	August 7	45.8	20.0	34.2	
	August 21	74.7	16.0	9.3	
	September 4	41.3	21.8	36.9	
	September 18	24.0	15.2	60.8	
	October 2	40.8	15.4	43.8	
	October 16	35.8	20.8	43.4	
	Mean		40.9	17.7	41.4
	95 percent confidence interval		31.3 - 50.6	15.7 - 19.6	31.0 - 51.8
	Tenmile	May 14	40.5	28.2	31.3
May 29		42.9	19.3	37.8	
June 10		30.4	11.5	58.1	
June 24		44.5	12.7	42.8	
July 22		61.4	17.5	21.1	
August 5		40.0	17.7	42.3	
August 19		32.5	16.2	51.3	
September 2		58.2	11.8	30.0	
September 16		50.3	17.0	32.7	
September 30		64.2	17.9	17.9	
October 14		8.1	6.2	85.7	
Mean			43.0	16.0	41.0
95-percent confidence interval			32.2 - 53.8	12.2 - 19.8	28.2 - 53.8

¹Surface to depth of 1.0 percent light

²Depth of 1.0 percent light to depth of 0.1 percent light

³Depth of 0.1 percent light to depth equal to depth of 0.1 percent light plus one-half of depth difference between 1.0 and 0.1 percent light depths.

a was contained within the euphotic zone and, thereby, provided evidence that phytoplankton was circulated out of the euphotic zone.

The importance of circulation of phytoplankton out of the euphotic zone in Lake Kooconusa was discussed by Woods (1981). He concluded that the reduction in phytoplankton photosynthesis resulting from reduced light intensities below the euphotic zone was an important factor in suppressing the reservoir's primary productivity to oligotrophic levels.

TROPHIC STATE OF LAKE KOOCANUSA

The trophic state of Lake Kooconusa can be categorized with one or more of the available trophic state indices. Most indices define discrete ranges for their categories of trophic state. A lake or reservoir generally can be placed within one of the following categories: Oligotrophic, mesotrophic, eutrophic, and hypereutrophic. Comparison of different trophic state indices, however, is often misleading because of the variety of variables used in their development. The use of different variables has arisen, in part, because three general approaches to trophic state determination are recognized. One approach considers nutrient supply as a stimulus to the water body. This approach uses vari-

ables such as nutrient concentrations within the water body or nutrient loads delivered to the water body from its drainage basin. The second approach uses the biological condition of the water body as an indicator of its trophic state. This approach uses variables such as primary productivity or chlorophyll *a* concentrations. The third approach combines the first two approaches to develop multivariate indices of trophic state.

The trophic state of Lake Kooconusa during 1972-80 was determined from data on primary productivity and loadings of total phosphorus and total nitrogen delivered into the reservoir. Primary productivity was used because it is a biological process that integrates the effects of numerous environmental variables controlling production of organic matter. Nutrient loadings were used because they have been emphasized in most studies of the aquatic environment of Lake Kooconusa, and Bonde and Bush (1975) used them to predict the trophic state of the reservoir prior to impoundment. The use of primary productivity and nutrient loadings for predicting trophic state was deemed appropriate in light of the conclusion drawn by Schindler (1978) that the relationship between nutrient supply and algal productivity has been well documented by numerous researchers.

A previous section of this report discussed primary productivity data that were used to classify the trophic state of Lake Kooconusa. The conclusion was drawn that Lake Kooconusa was oligotrophic from 1972-80 because its average daily areal primary productivity ranged from 63.6 to 105.5 mg·C·m⁻² (table 11) and, therefore, was clearly within the range of 50 to 300 mg·C·m⁻² that Wetzel (1975) considered oligotrophic.

The nutrient loading approach developed by Vollenweider (1968, 1975) was the other method used to classify the trophic state of the reservoir. Vollenweider (1968) was able to graphically separate 20 lakes into categories of oligotrophic, mesotrophic, and eutrophic by plotting logarithmic areal total phosphorus or total nitrogen loading of each lake against its logarithmic mean depth. Mean depth is the quotient of lake volume divided by surface area of the lake. Vollenweider also empirically determined a lower boundary line for permissible loading that quantified the maximum areal loading of total phosphorus or total nitrogen that a lake could receive and remain oligotrophic. Also determined was an upper boundary line for excessive loadings that would yield a eutrophic lake. Vollenweider (1975) modified his original model by adding hydraulic-residence time to allow for hydraulic effects in addition to the effects of nutrient loading and lake morphometry. The boundary lines for permissible and excessive loadings of total phosphorus were revised in the modified model; however, boundaries were not revised for total nitrogen

because the criteria needed to develop such boundary lines were incomplete (Rast and Lee, 1978).

Data needed to use Vollenweider's model include the yearly areal loading of total phosphorus or total nitrogen and the ratio of mean depth to hydraulic-residence time. Such data for Lake Kooconusa are listed in table 17; the data include 1972 through 1980 and the theoretical values for a reservoir containing 7.16 km³ and discharging an annual outflow of 10.65 km³, which is equal to its mean annual inflow. The theoretical nutrient loadings in table 17 are those reported by Bonde and Bush (1975) in their preimpoundment study of the reservoir.

The trophic state of Lake Kooconusa, based on application of Vollenweider's phosphorus loading model, is depicted in figure 8. The data in table 17 produced 10 points on figure 8; all 10 were within the eutrophic zone, but they were clustered into two groups. The theoretical value and those for 1972 through 1975 grouped higher into the eutrophic zone than the post-1975 group. The post-1975 group was lower than the pre-1975 group as a consequence of the 1976 reduction in phosphorus loading (table 4).

The relationship of areal total nitrogen loading and the ratio of mean depth to hydraulic residence time is depicted in figure 9 for Lake Kooconusa and 34 lakes analyzed by Rast and Lee (1978). The upper and lower boundary lines were not available for delineation of the three regions of trophic state, but the 10 points for Lake Kooconusa clearly occur within the cluster of eutrophic lakes. In contrast to figure 8, the 10 points were not split into two groups. The occurrence of the points in a single group was a consequence of the lack of reduction in nitrogen loading to the reservoir (table 5).

TABLE 17.—Required data for using Vollenweider^a nutrient loading models to determine the trophic state of Lake Kooconusa

Year	Mean depth, in meters	Hydraulic- residence time, in years	Ratio of mean depth to hydraulic- residence time, in meters per year	Annual areal loading, in grams per square meter	
				Total phosphorus	Total nitrogen
1972	23.2	0.14	165.7	15.7	61.7
1973	23.9	.33	72.4	20.7	41.1
1974	30.6	.29	105.5	11.9	32.4
1975	29.7	.41	72.4	10.5	22.5
1976	32.6	.38	85.8	3.8	22.6
1977	30.7	.50	61.4	2.8	19.3
1978	32.5	.48	67.7	3.7	21.3
1979	32.8	.62	52.9	2.9	19.1
1980	33.5	.58	57.8	2.9	22.1
Theoretical	38.4	.67	57.3	10.0 ^b	20.0 ^b

^aVollenweider (1975)

^bFrom Bonde and Bush (1975)

The eutrophic ranking that was derived for Lake Kooconusa from figures 8 and 9 conflicted with the oligotrophic ranking that the reservoir received for its values of primary productivity. The following discussion develops possible explanations for this discrepancy in trophic state ranking.

Two recent papers by Winter (1981a, 1981b) have emphasized measurement errors that are associated with water and chemical balances of lakes and reservoirs. Sufficient data were not available for Lake Kooconusa for estimating the magnitude of such errors for computations of annual areal loadings of total phosphorus and total nitrogen. However, the effect of these errors on the categorization of trophic state can be visualized with figures 8 and 9. If the annual areal loadings of total phosphorus in table 17 were assumed to be double their true values, the downward shifts in the 10 points that would be plotted on figure 8 for the true values still would be inadequate to place the reservoir into the oligotrophic zone. A similar conclusion was drawn for annual areal nitrogen loadings depicted in figure 9. Undoubtedly, errors existed in the water and chemical balances computed for both the reservoir and its various sources of inflow, but these errors did not fully account for the discrepancy in trophic state.

Only part of the nutrient loading that enters a lake or reservoir may become available to phytoplankton in the euphotic zone if the lake or reservoir experiences significant amounts of stratified interflow or underflow beneath the euphotic zone (Rast and Lee, 1978). Inflow to Lake Kooconusa during its annual filling phase was routed primarily as interflow or underflow (Woods, 1979). During the filling of Lake Kooconusa, the monthly loadings of total phosphorus and total nitrogen were at their largest for the year (Woods and Falter, 1982). Hence, some of the reservoir's nutrient supply may have been inaccessible to phytoplankton in the euphotic zone.

In addition to this hydrodynamic loss of nutrients, another part of the influent loading of total phosphorus may have been adsorbed and deposited by suspended sediment. Iskandar and Shukla (1981) concluded that Lake Kooconusa sediments functioned as a phosphorus sink.

The hydrodynamic and sedimentation processes within the reservoir help explain why potentially eutrophic nutrient loadings failed to stimulate primary productivity to eutrophic levels. However, the primary productivity of the reservoir was not solely controlled by nutrient supply; physical limnological processes contributed to the suppression of primary productivity to oligotrophic levels (Hobbie, 1976; Bush and Bonde, 1977; Woods, 1979, 1981). The essence of the reports by these authors is that the weak thermal structure of Lake




FIGURE 8.—Application of Lake Kooanusa data in table 17 to Vollenweider total phosphorus loading and mean depth/hydraulic-residence time diagram. Illustration modified from Rast and Lee (1978).

Kooanusa allowed circulation of phytoplankton out of an often shallow euphotic zone. Circulation into poor-light conditions reduced phytoplankton photosynthesis and, hence, annual primary productivity.

One can conclude that physical processes within Lake Kooanusa were more important in determining its trophic state, based on primary productivity, than was determination of its trophic state based on influent nutrient loadings. This conclusion supports a generalization expressed by Carlson (1979) that the use of nutrient loadings to characterize trophic state fails to consider the actual response of the lake; the nutrient loading

method only considers the lake's expected response to external stimuli.

CONCLUSIONS

Preimpoundment and postimpoundment studies of Lake Kooanusa have generated an extensive data base that has provided insight into the physical, chemical, and biological characteristics of the reservoir and its drainage basin. In these studies, the relationship of nutrient loadings, primary productivity, and trophic state has received much attention, largely because the preimpoundment studies reported large loadings of nitrogen




FIGURE 9.—Application of Lake Kooconusa data in table 17 to Vollenweider total nitrogen loading and mean depth/hydraulic-residence time diagram. Illustration modified from Rast and Lee (1978).

and phosphorus that might have caused eutrophication of Lake Kooconusa. During 1972–80, the reservoir was classifiable as eutrophic, based on the relationship of annual areal loadings of total nitrogen or total phosphorus to the ratio of mean depth and hydraulic-residence time. The eutrophic classification was not changed by the substantial reductions in total phosphorus loading that commenced in 1976 as a result of an upstream fertilizer plant instituting improved treatment of its effluent. However, the average daily areal primary productivity in Lake Kooconusa during 1972–80 characterized its trophic state as oligotrophic—in conflict with the eutrophic classification based on nutrient-loading data. This discrepancy of classification in trophic state is attributed to the failure of nutrient-loading models to adequately account for limnological processes within Lake Kooconusa, which affected the avail-

ability of influent nutrient loadings to phytoplankton.

Previous studies of limnological processes in Lake Kooconusa reported a weak thermal structure and, therefore, hypothesized that phytoplankton was circulated out of the euphotic zone. Such circulation was verified by a study of chlorophyll *a* distribution, which revealed that more than one-half of the reservoir's phytoplankton was located beneath the euphotic zone. Multiple regression analysis demonstrated that mean annual euphotic zone depth was an important predictor of annual areal primary productivity.

The relationship of daily values of primary productivity and concurrent values for physical, chemical, and biological variables was not studied in this project. Such an evaluation likely would yield valuable insight into the reasons why Lake Kooconusa was oligotrophic during 1972–80.

REFERENCES

- Bonde, T. J. H., 1979, Pre- and post-impoundment studies of the Kootenai River: Unpublished paper presented at the 109th annual meeting of the American Fisheries Society, West Yellowstone, Montana, September 14, 1979, 12 p.
- Bonde, T. J. H., and Bush, R. M., 1975, Kootenai River water quality investigations, Libby Dam preimpoundment study, 1967-1972: U.S. Army Corps of Engineers, Seattle District, 124 p.
- Brune, G. M., 1953, Trap efficiency of reservoirs: *American Geophysical Union Transactions*, v. 34, no. 3, p. 407-418.
- Bush, R. M., and Bonde, T. J. H., 1977, Relating water quality to the aquatic environment: Libby Dam-Lake Koocanusa project case study: Proceedings of a seminar on water quality data collection and management, Denver, Colorado, January 25-26, 1977, 10 p.
- Carlson, R. E., 1979, A review of the philosophy and construction of trophic state indices, in Maloney, T. E., ed., *Lake and reservoir classification systems*: U.S. Environmental Protection Agency, EPA-600/3-79-074, p. 1-52.
- Coffin, D. L., 1970, A preliminary evaluation of bank storage associated with Libby Reservoir in northwestern Montana: U.S. Geological Survey Water-Supply Paper 1899-L, 25 p.
- Crozier, R. J., and Leinweber, L. R., 1975, Libby Dam preimpoundment study: Nelson, British Columbia, British Columbia Pollution Control Branch, 166 p.
- Daley, R. J., Carmack, E. C., Gray, C. B. J., Pharo, C. H., Jasper, S., and Weigand, R. C., 1981, The effects of upstream impoundments on the limnology of Kootenay Lake, B. C.: Vancouver, British Columbia, Department of Environment, Inland Waters Directorate, 242 p.
- Dillon, P. J., and Rigler, F. H., 1974, A test of a simple nutrient budget model predicting the phosphorus concentration in lake water: *Journal of the Fisheries Research Board of Canada*, v. 32, p. 1777-1778.
- Draper, N. R., and Smith, H., 1966, *Applied regression analysis*: New York, John Wiley and Sons, 407 p.
- Fogg, G. E., 1974, Nitrogen fixation, in Stewart, W. D. P., ed., *Algal physiology and biochemistry*: Berkeley, University of California Press, p. 560-582.
- Golterman, H. L., 1975, *Physiological limnology*: Amsterdam, Elsevier Scientific Publishing Co., 489 p.
- Greeson, P. E., Ehlke, T. A., Irwin, G. A., Lium, B. W., and Slack, K. V., 1977, *Methods for collection and analysis of aquatic biological and microbiological samples*: U.S. Geological Survey Techniques of Water-Resources Investigations, Book 5, Chapter A4, 332 p.
- Hobbie, J. E., 1976, *Limnology of Lake Koocanusa, Montana*, in McKim, H. L., Gatto, L. W., Merry, C. J., Brockett, B. E., Bilello, M. A., Hobbie, J. E., and Brown, J., *Limnological Investigations: Lake Koocanusa, Montana, Part 2: Environmental Analyses in the Kootenai River Region, Montana*: Hanover, New Hampshire, U.S. Army Corps of Engineers, Cold Regions Research and Engineering Laboratory Special Report 76-13, p. 18-22.
- Iskandar, I. K., and Shukla, S. S., 1981, *Limnological investigations: Lake Koocanusa, Montana, Part 5: Phosphorus chemistry of sediments*: Hanover, New Hampshire, U.S. Army Corps of Engineers, Cold Regions Research and Engineering Laboratory Special Report 81-15, 9 p.
- Keeney, D. R., 1972, *The fate of nitrogen in aquatic ecosystems*: University of Wisconsin, Water Resources Center, Eutrophication Information Program, Literature Review No. 3, 59 p.
- Lee, G. F., 1970, *Factors affecting the transfer of materials between water and sediments*: University of Wisconsin, Water Resources Center, Eutrophication Information Program, Literature Review No. 1, 50 p.
- Likens, G. E., 1975, Primary productivity of inland aquatic ecosystems, in Leith, Helmut, and Whittaker, R. H., eds., *Primary productivity of the biosphere*: New York, Springer-Verlag Inc., p. 185-202.
- Rast, Walter, and Lee, G. F., 1978, Summary analysis of the North American (U.S. portion) OECD eutrophication project: nutrient loading-lake response relationships and trophic state indices: U.S. Environmental Protection Agency, EPA-600/3-78-008, 455 p.
- Reckhow, K. H., 1979, Quantitative techniques for the assessment of lake quality: U.S. Environmental Protection Agency, EPA-440/5-79-015, 146 p.
- Reckhow, K. H., Beaulac, M. N., and Simpson, J. T., 1980, Modeling phosphorus loading and lake response under uncertainty: A manual and compilation of export coefficients: U.S. Environmental Protection Agency, EPA-440/5-80-011, 214 p.
- Rigler, F. H., 1973, A dynamic view of the phosphorus cycle in lakes, in Griffith, E. J., Beeton, A., Spencer, J. M., and Mitchell, D. T., eds., *Environmental phosphorus handbook*: New York, John Wiley and Sons, p. 539-572.
- SAS Institute, Inc., 1979, *SAS user's guide*, 1979 edition: Cary, North Carolina, SAS Institute, Inc., 494 p.
- Schindler, D. W., 1978, Factors regulating phytoplankton production and standing crop in the world's freshwaters: *Limnology and Oceanography*, v. 23, p. 478-486.
- Slack, K. V., Averett, R. C., Greeson, P. E., and Lipscomb, R. G., 1973, *Methods for collection and analysis of aquatic biological and microbiological samples*: U.S. Geological Survey Techniques of Water-Resources Investigations, Book 5, Chapter A4, 165 p.
- U.S. Department of Commerce, issued annually, *Climatological data, Montana*.
- U.S. Environmental Protection Agency, 1977, *Report on Koocanusa Reservoir, Lincoln County, Montana, and British Columbia, Canada, EPA Region VIII: National Eutrophication Survey Working Paper No. 795*, 47 p.
- Vollenweider, R. A., 1968, Scientific fundamentals of the eutrophication of lakes and flowing waters, with particular reference to nitrogen and phosphorus as factors in eutrophication: Paris, Organization for Economic Cooperation and Development, Technical report DAS/CSI/68.27, 250 p.
- _____, 1975, Input-output models, with special reference to the phosphorus loading concept in limnology: *Schweizerische Zeitschrift fuer Hydrologie (Swiss Journal of Hydrology)* v. 37, p. 53-84.
- Water Resources Service, 1976, *Kootenay air and water quality study, Phase I, Water quality in Region 4, the Lower Kootenay River Basin*: Victoria, British Columbia, Department of Environment, 190 p.
- Wetzel, R. G., 1975, *Limnology*: Philadelphia, W. B. Saunders Co., 743 p.
- Winter, T. C., 1981a, Uncertainties in estimating the water balance of lakes: *Water Resources Bulletin*, v. 17, p. 82-115.
- _____, 1981b, Survey of errors for estimating water and chemical balances of lakes and reservoirs, in Stefan, H. G., ed., *Proceedings of a symposium on surface water impoundments*, v. 1: New York, American Society of Civil Engineers, p. 224-233.
- Woods, P. F., 1979, *Primary productivity in Lake Koocanusa, Montana*: Moscow, University of Idaho, Ph.D. dissertation, 112 p.
- _____, 1981, Physical limnological factors suppressing primary productivity in Lake Koocanusa, Montana, in Stefan, H. G., ed., *Proceedings of a symposium on surface water impoundments*, v. 2: New York, American Society of Civil Engineers, p. 1368-1377.
- Woods, P. F., and Falter, C. M., 1982, *Limnological investigations: Lake Koocanusa, Montana, Part 4: Factors controlling primary productivity*: Hanover, New Hampshire, U.S. Army Corps of Engineers, Cold Regions Research and Engineering Laboratory Special Report. In press.